



Rectangle and Trapezoid Approximations to Definite Integrals

Using the Document

RectangleTrapezoid.tns:

This file is used to graphically compare left, right, and midpoint rectangle approximations, and trapezoidal approximations to the area under a curve. Various examples are presented (using a single subinterval and single rectangle and/or trapezoid) to help determine when each of these approximations is an overestimate or underestimate, and to suggest which estimate produces the smallest error in estimating the area under a curve. Given certain characteristics of the curve, the graphs can be used to order the approximations. And the examples allow the user to describe how concavity affects the midpoint and trapezoidal approximations.

Suggested Applications and Extensions

Page 2.1

Use the slider arrow to view the shaded regions that represent the exact area under the curve and the estimates of the area under the curve.

1. Explain how each estimate region is constructed: left endpoint rectangle, right endpoint rectangle, and trapezoid.
2. Which of these three estimates is an overestimate for the exact area? Which of these is an underestimate? Which estimate do you think is the closest to the exact area?
3. Do you think that your answers to Problem 2 will be true for any function? Explain your reasoning.

Page 3.1

1. Use the slider to observe the estimates of the shaded area under the curve. Which of these three estimates is an overestimate for the exact area? Which is an underestimate? Are these results consistent with your answer to Problem 3 on Page 2.1? If so, explain why? If not, how would you adjust your prediction for overestimates and underestimates?
2. When will the trapezoid be an overestimate? When will it be an underestimate? When will it be a better estimate than the left endpoint rectangle and the right endpoint rectangle?

Pages 4.1 and 5.1

1. Use the slider to view the estimates of the shaded area under the curve, on both pages. Use your observations from Pages 2.1-5.1 to explain when a left endpoint rectangle is an underestimate or overestimate, when a right rectangle is an underestimate or overestimate, and when a trapezoid is an underestimate or overestimate.
2. Will a left rectangle ever provide the exact area under the curve? If so, how? If not, why not? Answer these same questions for a right rectangle and for a trapezoid.



Rectangle and Trapezoid Approximations to Definite Integrals

Page 6.1

1. Use the slider to view the exact area under the curve, the point on the curve used to construct the midpoint rectangle, and the midpoint rectangle. In your own words, explain how a midpoint rectangle is constructed. Why is it difficult to determine whether the area of a midpoint rectangle is an overestimate or underestimate?
2. Use the slider to select pivot. Click on pivot several times.
 - (a) Explain how the midpoint rectangle changes.
 - (b) As the top segment of the midpoint rectangle pivots, or rotates, how does the shaded area compare to the original midpoint rectangle? Justify your answer.
 - (c) Once the segment is done pivoting, a dashed line appears. What does the dashed line represent?
 - (d) Is the area of the midpoint rectangle an overestimate or an underestimate? Explain your reasoning.
3. In this case, which approach provides a more accurate estimate of the exact area under the curve: a midpoint rectangle or a trapezoid? Explain your reasoning.

Pages 7.1, 8.1, and 9.1

1. The functions presented on these pages are all nonnegative (over the subinterval considered), but may be increasing or decreasing, and may be concave up or concave down. Observe the estimates in each case and use your results from pages 3.1-9.1 to determine whether each approximation (L : left rectangle, R : right rectangle, M : midpoint rectangle, T : trapezoid) to the exact area under the curve is an underestimate or overestimate.
 - (a) The function is nonnegative, increasing, and concave up.
 - (b) The function is nonnegative, decreasing, and concave up.
 - (c) The function is nonnegative, increasing, and concave down.
 - (d) The function is nonnegative, decreasing and concave down.
2. How do your answers in Problem 1 change if *nonnegative* is replaced by *negative*? Explain.

In the next Problems, use your observations about a single subinterval to draw conclusions about the relationship among left, right, midpoint, and trapezoidal sums.

1. Suppose the function f is positive, continuous, increasing, and concave up on an interval $[a, b]$. Determine whether the inequality is true. Justify your answers.
 - (a) $L_n \leq M_n \leq R_n$
 - (b) $L_n \leq T_n \leq M_n$
 - (c) $M_n < \frac{L_n + R_n}{2}$
 - (d) $L_{2n} > L_n$



Rectangle and Trapezoid Approximations to Definite Integrals

2. Suppose the function f is positive, continuous, decreasing, and concave down on an interval $[a, b]$. Determine whether the inequality is true. Justify your answers.
- (a) $L_n \leq M_n \leq T_n \leq R_n$
 - (b) $L_n \geq M_n \geq T_n \geq R_n$
 - (c) $L_n \geq T_n \geq M_n \geq R_n$
 - (d) $L_n \geq R_n \geq M_n \geq T_n$
3. Let A be the area of the region bounded above by the graph of $y = e^{-x}$, below by the x -axis, on the left by the line $x = 0$, and on the right by the line $x = 8$.
- (a) List the values L_n , R_n , and A in increasing order. Justify your answer.
 - (b) Is T_n or M_n a better estimate of A ? Justify your answer.