

Name .	 
Class	 

## Problem 1 – Relating log functions with different bases

On page 1.3, you will see the graphs of two logarithmic functions with different bases:

 $f1(x) = \log_a(x) \text{ and } f2(x) = \log_b(x).$ 

- What are a and b? Trace the graphs to find out.
- What points on the graph are the best clues to the base of the logarithmic function?

Suppose we are interested in the sum of these two functions,  $(\mathbf{f1} + \mathbf{f2})(x) = \log_a(x) + \log_b(x)$ . How could we write this as a single logarithmic expression? We can't apply the properties of logarithms unless the logarithms have the same base. We need to rewrite the functions with the same base.

What does this mean? We want to find a function that is equal to **f1**, but has a log base *b* instead of log base *a*.

We could hope that there is a constant c that relates the two functions, like:  $c \cdot f1(x) = f2(x)$ .

Then, we would have  $\mathbf{f1}(x) = \frac{\mathbf{f2}(x)}{c} = \frac{1}{c} \cdot \log_b(x)$ , which is a logarithmic function with base b.

We can't be sure there is such a constant, but that doesn't have to stop us from looking for one.

• Solve  $c \cdot f1(x) = f2(x)$  for c and enter the result in f3 on page 1.4. What is c? View a function table to see.

## Problem 2 – A closer look at c

Is *c* always the same? Enter two new values of *a* and *b* in the gray cells on page 2.2. The value of *c* is calculated for you. Is it the same as in Problem 1? Record your results in the table below. Be sure to try some values of *a* and *b* such that one is a power of the other, like 2 and 8 or 3 and 9. Can you guess a formula for *c*?

а	f1( <i>x</i> )	b	f2(x)	С

## Problem 3 – Deriving the Change of Base Rule algebraically

We are convinced now that there is a constant that relates  $\log_a(x)$  to  $\log_b(x)$  and that the constant depends on the values of a and b. We may even have an idea what the constant is.

Time to use some algebra to find out for sure.

Two functions are equal if and only if their values are equal for every *x*-value in their domain. Let's pick a point (x, y) on the graph of **f1**. For this (x, y),  $\log_a(x) = y$ .

If we can write *y* in terms of logs base *b*, we will have our function.

•	Rewrite $log_a(x) = 1$	as an exponentia	I function.
•	TICWITC IOGALAL —	v as an expending	i iuiiouoii.

•	We want an ex	pression with	base b log.	so take le	og <sub>b</sub> of both sides
---	---------------	---------------	-------------	------------	-------------------------------

• Simplify using the properties of logs. Solve for *y*.

• What is *c*?

Check your equation for *c* against the value of *c* that you collected earlier. Enter it in the formula bar of Column D on page 2.2. When prompted, choose **Column Reference**, not Variable, for *a* and *b*. Is the equation correct?

You have found a formula for changing the base of a logarithm. To change a log base a expression to log base b, simply divide the expression by  $log_a(b)$ .

• Use this formula to find  $(\mathbf{f1} + \mathbf{f2})(x)$  if  $\mathbf{f1}(x) = \log_3(x)$  and  $\mathbf{f2}(x) = \log_{10}(x)$ .