# **Astroid**



## **Student Activity**

7 8 9 10 11 **12** 









### Introduction


How is the motion of a ladder sliding down a wall related to the motion of the valve on a bicycle wheel or to a popular amusement park ride? A little mathematical history, some parametric equations and calculus hold the key. Galileo first studied the cycloid in 1599. A cycloid describes the motion of the valve on a bicycle wheel. (Astroid TNS File: Page 1.2) Roemer (1674) put the cycloid into a 'spin' and developed the Astroid, not to be not to be confused with the celestial asteroid or Orbiter amusement park ride. The Astroid is a hypocycloid<sup>1</sup>.

# Cycloid

Open the TI-Nspire file: "Astroid". Navigate to page 1.2 of the file. Drag point A on the rim of the bicycle wheel and observe point P on the rim.

To see the trace (locus) of this point click on the 'show' button.

The size of the bicycle (wheel radius) can also be changed.




#### **Astroid**

Navigate to page 1.3.

Drag point L around the circle and watch the motion of point P.

To reveal the locus (path) of point P use the hide/show toggle.



#### Question: 1.

For each of the following questions: point L is moving counter-clockwise around the larger circle, the larger circle has a radius m measuring 8 units, the smaller circle has a radius of 2 units.

- a. Write down the parametric equations for the motion of the **centre** of the smaller circle.
- b. How many times does the smaller circle rotate as it travels around the inside of the larger circle?
- c. Consider the small circle as simply rotating on its own axis, located at the centre of the diagram; determine the parametric equations for the motion of point P.

  Make sure the equations account for the direction and number of rotations.

© Texas Instruments 2017. You may copy, communicate and modify this material for non-commercial educational purposes provided all acknowledgements associated with this material are maintained.

Author: P. Fox



<sup>1</sup> Hypo – means beneath, in this mathematical context it refers to a subset. So hypocycloid is a smaller set of the family of cycloids.

d. Combine your answers to part A and C and see if they follow the movement of Point P as it travels to form the Astroid.

#### Question: 2.

The parametric equations established in the previous question can be simplified.

- a. Show that  $\cos(3t) = 4\cos^3(t) 3\cos(t)$  and hence show that the parametric equation x(t) for the Astroid can be written as:  $x(t) = 8\cos^3(t)$
- b. Show that  $\sin(3t) = 3\sin(t) 4\sin^3(t)$  and hence show that the parametric equation for  $y(t) = 8\sin^3(t)$

#### Question: 3.

Use the parametric equations for the Astroid to show that an equivalent Cartesian equation can be expressed as:  $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$  and that for this specific case: a = 8.

#### Question: 4.

Use implicit differentiation techniques to determine the gradient of the Astroid at any point.

#### Question: 5.

Use the parametric equations for the Astroid to determine the gradient in terms of *t*.

#### Question: 6.

Show that the two expressions for the derivative (Cartesian and Parametric) are equivalent.

### Question: 7.

Determine the equation to the tangent for:  $t = \frac{\pi}{6}$ , the corresponding x and y intercepts, the length of the tangent joining the intercepts and the angle the tangent makes with the positive x axis.

#### Question: 8.

Determine the equation to the tangent for:  $t = \frac{\pi}{4}$ , the corresponding x and y intercepts, the length of the tangent joining the intercepts and the angle the tangent makes with the positive x axis.

#### Question: 9.

Determine the equation to the tangent for:  $t = \frac{\pi}{3}$ , the corresponding x and y intercepts, the length of the tangent joining the intercepts and the angle the tangent makes with the positive x axis.

#### Question: 10.

Comment on the angle the tangent makes with the x axis and your answer to question 5.

#### Question: 11.

Comment on the distance between the axes intercepts for each tangent equation.

#### Question: 12.

Use integral calculus to determine the distance that point P moves as it travels along the length of the Astroid.



# **Falling Ladder**

Navigate to page 2.1.

Drag point P around the ground and watch the motion of the ladder.


From the menu select:


# **Geometry > Construction > Locus**

Select the ladder followed by point P.

Navigate to page 2.2

Drag point P along the x axis. Use the toggle to show a trace of the ladder's position both above and below the x axis.





# **Exploration**

Explore the motion of the ladder. Show that the envelope traced out by the movement of the ladder forms an Astroid?