



Math Objectives

- Students will discover that the zeros of the linear factors are the zeros of the polynomial function.
- Students will discover that the real zeros of a polynomial function are the zeros of its linear factors.
- Students will determine the linear factors of a quadratic function.
- Students will connect the algebraic representation to the geometric representation.
- Students will see the effects of a double and/or triple root on the graph of a cubic function.
- Students will see the effects of the leading coefficient on a cubic function.
- Students will look for and make use of structure (CCSS Mathematical Practice).
- Students will look for and express regularity in repeated reasoning (CCSS Mathematical Practice).

Vocabulary

- zeros
- double or triple root
- leading coefficient

About the Lesson

- This lesson merges graphical and algebraic representations of a polynomial function and its linear factors.
- As a result, students will:
 - Manipulate the parameters of the linear functions and observe the resulting changes in the polynomial function.
 - Find the zeros of the polynomial equations by finding the zeros of the linear factors.




TI-Nspire™ Navigator™

- Use Class Capture to examine patterns that emerge.
- Use TI-Nspire Teacher software or Live Presenter to review student documents and discuss examples as a class.
- Use Quick Poll to assess student understanding.

Activity Materials

- Compatible TI Technologies:  TI-Nspire™ CX Handhelds,  TI-Nspire™ Apps for iPad®,  TI-Nspire™ Software



Tech Tips:

- This activity includes screen captures taken from the TI-Nspire CX handheld. It is also appropriate for use with the TI-Nspire family of products including TI-Nspire software and TI-Nspire App. Slight variations to these directions may be required if using other technologies besides the handheld.
- Watch for additional Tech Tips throughout the activity for the specific technology you are using.
- Access free tutorials at <http://education.ti.com/calculators/pd/US/Online-Learning/Tutorials>

Lesson Files:

Student Activity

- Exploring_Polynomials_Factors_Roots_and_Zeros_Student.pdf
- Exploring_Polynomials_Factors_Roots_and_Zeros_Student.doc


TI-Nspire document

- Exploring_Polynomials_Factors_Roots_and_Zeros.tns



Discussion Points and Possible Answers



Tech Tip: If students experience difficulty clicking a slider, check to make sure that they have moved the cursor over the slider and have them press  to change the value of the slider.



Tech Tip: Tap on the arrows to change the values of the slider.




TI-Nspire Navigator Opportunity: *Live Presenter* or *Class Capture*

See Note 1 at the end of this lesson.

Move to page 1.2.

1. Using the sliders, set $y_1 = 1x + 1$ and $y_2 = 1x - 2$. Observe that the graph of $y_1 = 1x + 1$ appears to cross the x-axis at $x = -1$. When $x = -1$, $y_1 = 0$ because $-1 + 1 = 0$. $x = -1$ is called a *zero* or *root* of the function $y_1 = 1x + 1$.



- a. Where does the graph of $y_2 = 1x - 2$ appear to cross the x-axis?

Answer: $x = 2$

- b. Write a simple equation to verify that this value of x is a zero of y_2 .

Answer: $1(2) - 2 = 0$

Teacher Tip: A zero of a function is an input value for which the function value is zero. Thus, if $x = 2$ is a zero of the function, then $f(2) = 0$ and the point $(2, 0)$ is on the graph of the function.

- c. When $y_1 = 1x + 1$ and $y_2 = 1x - 2$, what is the function y_3 ?

Answer: $y_3 = x^2 - x - 2$

- d. The graph of y_3 is a parabola. How many times does the graph of y_3 cross the x-axis?

Answer: The graph crosses the x-axis twice.



- e. What are the zeros of y_3 ?

Answer: $x = -1$ and $x = 2$

- f. Factor y_3 .

Answer: The factors of $x^2 - x - 2$ are $(x + 1)$ and $(x - 2)$.

Teacher Tip: This activity is assuming that the method of factoring a quadratic has already been completed. Teachers may need to do some reviewing of factoring at this point.



Tech Tip: To change the slider settings, press and hold on an arrow. Select “Settings.” Then change any values in the Settings menu.

- g. Given the information below, use the sliders to fill in the rest of the table:

Answer: Completed table is below. Answers may vary because students may choose to factor completely. It is acceptable for linear factors not to be completely factored.

y_1	y_2	Zeros of y_1 y_2		y_3	Zeros of y_3	Factors of y_3
$(x + 4)$	$(x + 3)$	-4	-3	$x^2 + 7x + 12$	-4 and -3	$(x + 4)(x + 3)$
$(2x - 4)$	$(x + 2)$	2	-2	$2x^2 + 0x - 8$	2 and -2	$(2x - 4)(x + 2)$
$(x - 5)$	$(-1x - 2)$	5	-2	$-1x^2 + 3x + 10$	5 and -2	$(x - 5)(-1x - 2)$
$(3x + 3)$	$(x + 4)$	-1	-4	$3x^2 + 15x + 12$	-1 and -4	$(3x + 3)(x + 4)$
$(x + 1)$	$(x - 4)$	-1	4	$x^2 - 3x - 4$	-1 and 4	$(x + 1)(x - 4)$
$(2x + 4)$	$(3x - 3)$	-2	1	$6x^2 + 6x - 12$	-2 and 1	$(2x + 4)(3x - 3)$

- h. Write a conjecture about the relationship between the zeros of the linear functions and the zeros of the quadratic function.

Answer: The zeros of the linear functions are the zeros of the quadratic function.

Teacher Tip: Some of the polynomials are not fully factored. This is a topic you might choose to explore with the students.



- i. How do the factors of the quadratic equation relate to the zeros of the function?

Answer: The factors of the quadratics are the same linear functions that multiply together to make the quadratic and therefore can be solved to find the zeros or x-intercepts of the quadratic function. If a polynomial can be factored, factoring is one strategy for finding the real solutions of a polynomial equation.

Teacher Tip: If students haven't solved quadratics by factoring, this would be a good time to discuss the concept.


Move to page 2.2.

2. Use the sliders to make $f1 = 1x + 4$, $f2 = 1x + 2$, and $f3 = 1x - 1$.

Observe that the graphs of each appear to cross the x-axis at -4 , -2 , and 1 , respectively.

- a. Verify algebraically that each is a zero of each linear function.

Answer: $1(-4) + 4 = 0$, $1(-2) + 2 = 0$, $1(1) - 1 = 0$



- b. When $f1 = 1x + 4$, $f2 = 1x + 2$, and $f3 = x - 1$, what is $f4$?

Answer: $f4 = x^3 + 5x^2 + 2x - 8$

- c. How many times does $f4$ cross the x-axis and where?

Answer: Three times: at -4 , -2 , and 1

- d. Show the multiplication of the factors of $f1$, $f2$, and $f3$ to equal $f4$.

Answer: $(x + 4)(x + 2) = x^2 + 6x + 8$, then $(x^2 + 6x + 8)(x - 1) = x^3 + 5x^2 + 2x - 8$

Teacher Tip: If necessary, review multiplication of polynomial expressions by distributing each term in the first parentheses by every term in the second parentheses.


- e. Try other slider values and make a conjecture about the relationship between the zeros of the linear equations and the zeros of the cubic function.

Answer: The zeros of the linear functions are the zeros of the cubic function.



3. Use the sliders to make $f1 = x + 4$, $f2 = x + 2$, and $f3 = x + 2$.
- a. How has the graph changed? The value -2 is called a double root.


Sample answer: Answers may vary, but students should say something similar to the fact that the graph no longer crosses the x -axis in three places, but appears to “bounce back up” at -2 . It still crosses at -4 . The new equation is $f4 = x^3 + 8x^2 + 20x + 16$.



Teacher Tip: By changing one of the factors to $0x$ and then getting a parabola, students might see that this point is the vertex of the quadratic, as seen previously.


- b. Change $f1 = 1x + 2$. How has the graph changed?

Sample answer: Answers may vary, but the graph appears to flatten out at -2 . The value -2 is a triple root of $f4 = x^3 + 6x^2 + 12x + 8$.




4. Use the sliders to make $f1 = 3x - 3$, $f2 = x + 1$, and $f3 = x - 2$.
- a. Observe the graph and identify the zeros. What is $f4$?

Answer: The zeros are 1 , -1 , and 2 , and $f4 = 3x^3 - 6x^2 - 3x + 6$.



- b. Now change the sliders to make $f1 = x - 1$, $f2 = x + 1$, and $f3 = x - 2$. Observe the graph. What are the zeros? What is $f4$?

Answer: The zeros are 1 , -1 , and 2 , and $f4 = x^3 - 2x^2 - x + 2$.





- c. Identify similarities and differences between the sets of equations in part a and part b.

Answer: The zeros of both functions are the same. The graph rises or falls differently between the two graphs. The second function is the first function multiplied by a factor of 3. The leading coefficient causes a vertical dilation (factor) of 3. Each factor can be used to find the zeros or x-intercepts of the functions or to find the roots of the corresponding equations.

Teacher Tip: By changing one of the factors to $0x$ and then getting a parabola, students might see that this point is the vertex of the quadratic, as seen previously.



TI-Nspire Navigator Opportunity: Quick Poll

See Note 2 at the end of this lesson.

Wrap Up

Upon completion of the discussion, the teacher should ensure that students understand:

- How to use a graph to find possible linear factors of a quadratic function.
- The connections between the algebraic and graphical representations of a quadratic and cubic function and its factors.
- That the zeros of the linear factors of a polynomial function and the zeros of the polynomial function are the same.
- That the zeros of a polynomial function are the same as the zeros of its linear factors.
- How a double or triple root of a polynomial function affects the graph.
- The effects of the leading coefficient on a cubic function.

Assessment

1. Given zeros of -4.5 , -1 , and 2 , find a possible cubic equation. Is your answer unique? Explain.
2. Given that $(x + 5)$ and $(2x - 1)$ are the only factors of a cubic polynomial, find a possible cubic equation. Is your answer unique? Explain.



TI-Nspire Navigator

Note 1

Entire Lesson, *Live Presenter* or *Class Capture*: If students experience difficulty with the sliders, use *Class Capture* or *Live Presenter* with TI-Nspire Navigator to demonstrate to the entire class.

Note 2

End of Lesson, *Quick Polls*: The following *Quick Poll* questions can be given at the conclusion of the lesson. You can save the results and show a class analysis at the start of the next class to discuss possible misunderstandings students may have.

1. Given zeros of -5 , 1 , and 3 , a possible cubic equation is:

- a. $y = (x - 5)(x + 1)(x + 3)$
- b. $y = (x + 5)(x - 1)(x - 3)$
- c. $y = (x - 5)(x - 1)(x + 3)$
- d. $y = (x + 5)(x - 1)(x + 3)$

Answer: b

2. The zeros of $y = x(x + 4)(x - 2)$ are:

- a. $1, -4, 2$
- b. $0, 4, -2$
- c. $1, 4, -2$
- d. $0, -4, 2$

Answer: d

3. A cubic equation has a root at -6 and a double root at 4 . The factors of the equation are:

- a. $(x + 6)$, $(x + 4)$, and $(x - 4)$
- b. $(x - 4)$, $(x + 6)$, and $2(x - 4)$
- c. $(x + 6)$, $(x - 4)$, and $(x - 4)$
- d. $(x - 6)$, $(x + 4)$, and $(x + 4)$

Answer: c